PRODUCT INFORMATION

Product name: EZH1 antibody

Product type: Primary antibodies

Description: Rabbit polyclonal to EZH1

Immunogen: 3 synthetic peptides (human) conjugated to KLH

Reacts with: Human, Mouse

Tested applications: ELISA, WB

GENE INFORMATION

Gene Symbol: EZH1

Gene Name: enhancer of zeste homolog 1 (Drosophila)

Ensembl ID: ENSG00000108799

Entrez GeneID: 2145 Omim ID: 601674 Swiss-Prot: Q92800

Molecular weight of EZH1: 85.2kDa

Function: Polycomb group (PcG) protein. Catalytic subunit of the PRC2/EED-EZH1 complex, which methylates 'Lys-27' of histone H3, leading to transcriptional repression of the affected target gene. Able to mono-, di- and trimethylate 'Lys-27' of histone H3 to form H3K27me1, H3K27me2 and H3K27me3, respectively. Required for embryonic stem cell derivation and self-renewal, suggesting that it is involved in safeguarding embryonic stem cell identity. Compared to EZH1-containing complexes, it is less abundant in embryonic stem cells, has weak methyltransferase activity and plays a less critical role in forming H3K27me3, which is required for embryonic stem cell identity and proper differentiation

Expected subcellular localization: Nucleus. Note: Colocalizes with trimethylated 'Lys-27' of histone H3

Summary: EZH1 is a component of a noncanonical Polycomb repressive complex-2 (PRC2) that mediates methylation of histone H3 (see MIM 602812) lys27 (H3K27) and functions in the maintenance of embryonic stem cell pluripotency and plasticity (Shen et al., 2008 [PubMed 19026780]).[supplied by OMIM, Mar 2009]

APPLICATION NOTE

Recommended dilution:

- ELISA: Antibody specificity was verified by direct ELISA against the 3 immunogen peptides. A minimum titer of 1/25000 is determined. Appropriate specificity controls were run.
- WB: 1/5000.

Optimal dilutions/concentration should be determined by the end user.

Raised in: Rabbit

Clonality: Polyclonal

Isotype: IgG

Purity: Purified Antibody

Storage buffer: 0.5 X PBS, 50% glycerol containing a final concentration of 0.1% BSA and

0.01% Thimerosal. Form: Liquid


Storage instruction: Store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

The purified antibody ENSG00000108799 has been tested at 1/5000 on uninduced (negative control) and induced culture of E.coli (one shot Top10 competent cells).

An anti-thio has been tested at 1/5000 on induced culture of E.coli (one shot Top10 competent cells) as a positive control.

Plasmid name: pBAD-DEST49.

Molecular weight of EZH1: 99.3kDa (85.3kDa + another 14kDa for the tag).



Gel concentration: 10%

Blocking: in 5% non-fat milk-PBST solution

1st Antibody: The antibodies are diluted in blocking buffer.

- Dilute the purified antibody ENSG00000108799 at 1: 5000
- Dilute the anti-thio at 1:5000

60 minutes of incubation

2nd Antibody: The antibody is diluted in blocking buffer.

• Dilute the anti-Rabbit IgG HRP conjugated at 1/10000 60 minutes of incubation